

## Presence of *Radix lagotis* (Schrank, 1803) (Gastropoda: Lymnaeidae) in the NE Iberian Peninsula

Joaquín López Soriano<sup>1,\*</sup>, Sergio Quiñonero Salgado<sup>2</sup> & Peter Glöer<sup>3</sup>

<sup>1</sup>Vall d'Hebron Institut de Recerca (VHIR), Passeig Vall d'Hebron 119-129, 08035 Barcelona, Spain; <sup>2</sup>Associació Catalana de Malacologia, Museu Blau, Plaça Leonardo da Vinci 4-5, 08019 Barcelona; <sup>3</sup>Biodiversity Research Laboratory, Schulstrasse 3, D-25491 Hetlingen, Germany.

Rebut el 30 d'agost de 2016

Acceptat el 17 de setembre de 2016

© Associació Catalana de Malacologia (2016)

*Radix lagotis* (Schrank, 1803) is a lymnaeid species that inhabits ponds, rivers and lakes in the Palaearctic region. It has been reported from many European countries, particularly in the Danube basin (Germany, Austria, Bulgaria, Czech Republic, Romania and Slovakia; Bank *et al.*, 2006 in von Proschwitz, 2011; Cioboiu, 2006), and also far in the east (Russia and Kyrgyzstan; Vinarski *et al.*, 2007; Glöer *et al.*, 2014). However, it had never been found in the Iberian Peninsula. It has a small (about 15–25 mm), thin, shiny shell with 4–5 whorls, deep sutures, and a moderately inflated last whorl; the columella is usually curved and weakly folded, and the umbilicus is very narrow (Glöer, 2002). Here we report the first citation of this species for the Iberian malacofauna, based on several finds from some tributaries of the Ebro River (Figures 1–3):

- Algars River at Arnes (la Terra Alta, Tarragona) [31T BF6933], 452 m; 16/4/2016 JLS & SQS *leg.* Alive specimens collected in shallow water among vegetation. This river marks the border between Catalonia and Aragon at this locality, so the species should be considered present in both autonomous communities.
- La Canaleta River at Horta de Sant Joan (la Terra Alta, Tarragona) [31T BF7432], 518 m; 16/4/2016 JLS & SQS *leg.* Alive specimens collected in shallow water among vegetation.
- Els Estrets River at Horta de Sant Joan (la Terra Alta, Tarragona) [31T BF7328], 551 m; 16/4/2016 JLS & SQS *leg.* Alive specimens taken in shallow water among vegetation.

*Radix lagotis* can only be confused with two other species present in the Iberian Peninsula: *Radix balthica* (Linnaeus, 1758) and *Radix labiata* (Rossmäessler, 1835). The former is present in Catalonia, including the Ebro River and Delta (Alba *et al.*, 2011; Quiñonero Salgado & López Soriano, 2014). *Radix labiata*, in turn, has been reported from high mountain habitats at Santa Marina de Valdeón (León), in the Picos de Europa lakes and marshes (Schniebs *et al.*, 2013), and recently also from mountain rivers in Aragon, after anatomical revision of some specimens (Quiñonero Salgado *et al.*, 2016). Shell morphology does not seem to be a definitive diagnostic criterion for the differentiation of the three species mentioned above (given their overall similarity and high degree of shape plasticity), but DNA analyses have confirmed their status as different species (Bargues *et al.*, 2001; Schniebs *et al.*, 2013). Moreover, a comparison of the bursa duct of *R. lagotis* with that of its closest relatives, *R. balthica* and *R. labiata*, enables a reliable distinction of these taxa (Schniebs *et al.*, 2011, 2013; Glöer, 2015). This has been considered as the best anatomical distinguishing character between these closely-related species (Schniebs *et al.*, 2013). The bursa duct of *R. lagotis* is longer than that of *R. labiata* (reaching about two-thirds of the bursa length), whereas it is much shorter in *R. balthica*. The anatomy of the



**Figure 1.** Map of Catalonia (within the Iberian Peninsula) showing the location of the new finds of *Radix lagotis* reported herein.

specimens studied by us from Catalonia (Figure 3) perfectly matches that reported for *R. lagotis* by Schniebs *et al.* (2013).

All our finds of *R. lagotis* in Catalonia are located in tributaries of the Ebro River within the natural park of Els Ports, a protected area with pristine waters and calcareous soils. Therefore, it is conceivable that this species may be common and widespread in the water courses of these mountains, although extensive searches would be necessary to verify this point. This species has been described as an intermediate host for the nasal bird schistosome *Trichobilharzia regenti*, a parasite which is also responsible of some cercarial dermatitis in humans (Skála *et al.*, 2014), although we are not aware of such cases in this part of the territory.


After the finds reported here, three species of *Radix* are known in Catalonia: *Radix auricularia* (Linnaeus, 1758), *R. balthica*, and *R. lagotis*. Two other species of the genus are also present in Spain (Glöer & Beckmann, 2007; Schniebs *et al.*, 2013): *R. labiata*, at least in N and NW Spain; and *Radix lilli* Glöer & Beckmann, 2007, endemic of the Balearic Islands. The results reported herein, added to the report (based on anatomical data) of *R. labiata* and *Stagnicola fuscus* (Pfeiffer, 1821) in Spain (Quiñonero Salgado *et al.*, 2016; López Soriano *et al.*, 2016), suggest that a revision of the previous reports of lymnaeid species in Spain is urgently needed, as most of their citations were only based on conchological traits—which are not quite reliable for members of this family without an anatomic confirmation.

### Acknowledgements


We want to thank Marco Pla for field trips and information about the populations studied in this article, as well as Jordi Corbella, Josep Quintana and David M. Alba for critical review of the manuscript.

\* Autor corresposnal.

Adreça electrònica: [qlopez@yahoo.com](mailto:qlopez@yahoo.com)



**Figure 2.** A, Several views of the shell of a representative specimen of *Radix lagotis* from Horta de Sant Joan (Tarragona); from left to right: dorsal, lateral, and ventral views. B, Alive specimen of *R. lagotis* in its shallow-water habitat at Algars River (Arnes, Tarragona). C, Habitat of *R. lagotis* at Algars Rivers, where many specimens were found.



**Figure 3.** Gonad morphology of *Radix lagotis* from Horta de Sant Joan (Tarragona). Abbreviations: bc = bursa copulatrix; bd = bursa duct; cp = corpus pyriforme; pvd = provaginal duct; prp = praeputium; pht = phalotheca; vd = vas deferens; pg = prostate gland.

## References

- Alba, D.M., Tarruella, A., Prats, L., Guillén, G. & Corbella, J. (2011). Nova llista actualitzada dels mol·luscos continentals de Catalunya. *Spira* 4, 39–69.
- Bank, R., von Proschwitz, T. & Falkner, G. (2006). Unpublished manuscript of the mollusca section of the Fauna Europea web-site. <http://www.faunaeur.org>
- Bargues, M.D., Vigo, M., Horak, P., Dvorak, J., Patzner, R.A., Pointier, J.P., Jackiewicz, M., Meier-Brook, C. & Mas-Coma, S. (2001). European Lymnaeidae (Mollusca: Gastropoda), intermediate hosts of trematodiasis, based on nuclear ribosomal DNA ITS-2 sequences. *Inf. Gen. Evol.* 16, 1–23.
- Cioboiu, O. (2006). Diversity of Gastropoda in the Romanian sector of the Danube lower hydrographic basin. In: *Proceedings 36th International Conference of International Association of Danube Research*, p. 230–235. Austrian Committee Danube Research / IAD, Vienna.
- Glöer, P. (2002). *Süßwassergastropoden Nord- und Mitteleuropas*. Conchbooks, Hackenheim.
- Glöer, P. (2015). *Süßwassermollusken. Ein Bestimmungsschlüssel für die Muscheln und Schnecken im Süßwasser der Bundesrepublik Deutschland*. Deutscher Jugendbund für Naturbeobachtung, Hamburg.
- Glöer, P. & Beckmann, K.-H. (2007). *Radix lilli* n. sp. und drei neue *Bithynia*-Arten von den Balearen (Gastropoda: Bithyniidae, Lymnaeidae). In: Beckmann, K.-H., *Die Land- und Süßwassermollusken der Balearischen Inseln*, 163–170. Conchbooks, Hackenheim.
- Glöer, P., Boeters, H.D. & Pešić, V. (2014). Freshwater molluscs of Kyrgyzstan with description of one new genus and species (Mollusca: Gastropoda). *Folia Malacol.* 22, 73–81.
- López Soriano, J., Quiñonero Salgado, S. & Glöer, P. (2016). The genus *Stagnicola* Jeffreys, 1830 in Catalonia. *Spira* 6, 81–83.
- Quiñonero Salgado, S. & López Soriano, J. (2014). Moluscos continentales del delta del Ebro (Cataluña, España). *Spira* 5, 121–132.
- Quiñonero Salgado, S., López Soriano, J., Glöer, P. (2016). First citation for *Radix labiata* (Rossmässler, 1835) (Gastropoda: Lymnaeidae) in Aragon (NE Spain). *Spira* 6, 85–86.
- Schniebs, K., Glöer, P., Vinarski, M.V. & Hundsdoerfer, A.K. (2011). Intraspecific morphological and genetic variability in *Radix balthica* (Linnaeus, 1758) (Gastropoda: Basommatophora: Lymnaeidae) with morphological comparison to other European *Radix* species. *J. Conchol.* 40, 657–678.
- Schniebs, K., Glöer, P., Vinarski, M.V. & Hundsdoerfer, A.K. (2013). Intraspecific morphological and genetic variability in the European freshwater snail *Radix labiata* (Rossmässler, 1835) (Gastropoda: Basommatophora: Lymnaeidae). *Contrib. Zool.* 82, 55–68.
- Skála, V., Černíková, A., Jindrová, Z., Kašný, M., Vostrý, M., Walker, A.J. & Horák, P. (2014). Influence of *Trichobilharzia regenti* (Digenea: Schistosomatidae) on the defence activity of *Radix lagotis* (Lymnaeidae) haemocytes. *PLoS ONE* 9, e111696.
- Vinarski, M.V., Karimov, A.V., Grebennikov, M.E. & Lazutkina, E.A. (2007). Aquatic gastropods of the Ilmeny State Reserve (Southern Urals, Russia). *Tentacle* 15, 8–10.
- von Proschwitz, T. (2011). *Radix lagotis*. In: *The IUCN Red List of Threatened Species 2011*, e.T155502A4788468. <http://dx.doi.org/10.2305/IUCN.UK.2011-2.RLTS.T155502A4788468.en> [Accessed 17/9/2016]